

Prozap[®] Annihilator-XP

EMULSIFIABLE CONCENTRATE

Multi-purpose insecticide for use on ornamentals, stored products and livestock areas.

Use alone as a clean-up spray or a pre-harvest spray.

GUARANTEE:

Pyrethrins6.00%
 *Piperonyl Butoxide60.00%

*Equivalent to 48.00% (butylcarbityl) (6-propylpiperonyl) ether and 12.00% related compounds.

Registration No. 31943 PCP Act

EPA Est. No. 47000-IA-01

READ THE LABEL BEFORE USING

COMMERCIAL

SKIN IRRITANT

POTENTIAL SKIN SENSITIZER

CAUTION [SKULL BONES] POISON

KEEP OUT OF REACH OF CHILDREN

FIRST AID	
If swallowed:	Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration number with you when seeking medical attention.
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.
Take container, label or product name and Pest Control Product Registration number with you when seeking medical attention.	

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Avoid contact

with skin or eyes. Wash with soap and warm water after use. Avoid contamination of food and feedstuffs. Do not apply space spray while food processing is underway. Cover or remove all food processing surfaces. Thoroughly wash all food processing surfaces before reuse. Remove all pets and birds and cover fish aquariums before spraying. This product is toxic to fish. Do not apply directly to any body of water by cleaning of equipment or disposal of wastes. Do not use or store near heat or open flame. When mixing, loading and applying, wear a chemical resistant apron over long sleeved shirt and long pants, chemical resistant gloves, and socks plus chemical resistant foot wear.

TOXICOLOGICAL INFORMATION: Contains petroleum distillate. Vomiting may cause aspiration pneumonia. Treat symptomatically.

DIRECTIONS FOR USE:

Multi-purpose insecticide for use on ornamentals, stored products and livestock areas. Use alone as a clean-up spray or a pre-harvest spray.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The areas being treated must be vacated by unprotected persons. Do not enter treated areas until sprays have dried.

USED ALONE: This concentrate can be used also as a clean-up or a pre-harvest spray where other materials cannot be used because of residue restrictions. This product may be used up to and including day of harvest. Contains natural Pyrethrins.

USED DILUTED: Apply 60 to 480 milliliters per .4 hectares and repeat if required to maintain effective control. Use in sufficient water for thorough coverage of upper and lower leaf surfaces (maximum of 380 L of water per .4 hectares) unless otherwise noted. It is recommended that the final spray mix be buffered to a PH of 5.5-7.0.

Do not apply this product through any type of irrigation system.

This concentrate is relatively non-toxic to honey bees. To avoid possible harm to honey bees, it is advisable to apply in the early morning or late evening hours.

ORNAMENTALS: Such as African Violet Aster, Azalea, Begonia, Calceolaria, Calendula, Calla, Camellia, Carnation, Cineraria, Chrysanthemum, Cypress, Daffodil, Dahlia, Dogwood, Elm Eucalyptus, Fern, Ficus, Geranium, Gladiolus, Gypsophila, Holly Juniper, Lily, Marigold, Oak, Peony, Petunia, Philodendron, Pine, Roses, Snapdragon, Sweet Pea, Tulips, Viburnum, Wandering Jew, yew, Zinnia.

FOR THE CONTROL OF INSECTS:

Such as Ants, Aphids, Armyworms, Asparagus Beetles, Blister Beetles, Cabbage Looper, Caterpillars, Cockroaches, 12-spotted cucumber beetle, Colorado Potato Beetles, Corn Earworm, Crickets, Cross-striped Cabbageworm, Cucumber Beetles, Deer Fly, Diamondback larvae, Fireworms, Flea Beetles, Fruit Flies, Fruittree Leafroller, Grape Leafhopper, Green Peach Aphids, Greenhouse Thrips, Gypsy Moth (adults & larvae), Harlequin bug, Heliothis sp., Hornets, Horn Fly, Horse Fly, House Fly, Imported Cabbageworm, Leafhopper, Leafroller, Leaf tiers, Lice, Mexican Bean Beetle, Mosquitoes, Potato Leafhopper, Psyllids, Silverfish, Skippers, Stable Fly, Stink Bugs, Tabanids, Thrips, Vinegar Flies, Wasps, Webworms, Whiteflies and Yellowjackets.

FOR USE AROUND HOMES AND OTHER BUILDINGS: Spray grassy areas around yard borders thoroughly, contacting as many insects as possible on upper and lower leaf surfaces to control ticks that may carry Lyme disease. Spraying should be done when the insects first appear. Do not wait until the plants are heavily infested.

FOR USE IN FOOD AND NONFOOD AREAS OF FOOD PROCESSING PLANTS, INDUSTRIAL INSTALLATION, BAKERIES, RICE AND WHEAT MILLS, RESTUARANTS, TOBACCO WAREHOUSES, GRAIN ELEVATORS AND WAREHOUSES: To kill flying insects such as fruit flies, house flies, hornets, wasps, grain moths, gnats, mosquitoes and skipper flies, dilute this concentrate at the rate of 1 part with 29 parts water (.95L with 28.50L water) up to 1 part to 11 parts water (.95L with 11.4L water). Use at the rate of 15 to 30 milliliters of dilute spray per 28.30 cu. meters of space. Direct the space treatment upward and whenever practical, keep doors product in food processing or food handling establishments should be confined to time periods when the plant is not in operation. Food should be covered or removed during treatment.

Where oil residues are not undesirable, this product can be diluted at the rate of 1 part to 29 parts up to 1 part to 11 parts in deodorized base oil instead of water and applied as a space spray with any good type applicator such as mechanical or ULV fogger capable of producing particles of aerosol size.

FOR USE IN STORAGE SITES: This concentrate can be used in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment as a clean-up prior to using them for storage. In mills and elevators, all grain infested accumulations should be removed from the bin hoppers. All storage areas and conveying equipment should be thoroughly cleaned by sweeping out the waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and door frames with special attention to material lodged in the cracks and crevices. All of the debris should be removed and burned to kill eggs and insects that might be present.

For farms, particular attention should be given to cleaning up around the used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. Newly harvested grain should not be placed in the same bin with carry-over grain and all carry-over grain stocks that are not treated with grain protectant should be fumigated. These

cleaning operations should be done within two or three weeks before harvest.

After above sanitation measures have been employed, spray all areas prior to use for storage with 1 part to 29 parts water (.95L with 28.50L water) up to 1 part to 11 parts (.95L with 11.4L water). Apply at the rate of 3.8L per 70 sq. meters on walls, floors, ceilings and partition boards of bins, paying particular attention to forcing the spray into all cracks and crevices.

FOR USE IN LIVESTOCK AREAS:

1. To control poultry lice, using a dilution of 60 to 90 milliliters of concentrate per 3.8L of water spray roosts, walls and nests or cages thoroughly.
2. For control of bedbugs and mites in poultry houses, dilute at the rate of 60 to 90 milliliters per 3.8L of water and spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.

FOR USE IN BARN, DAIRIES, MILKING PARLORS, MILKING ROOMS AND POULTRY HOUSES: To control flies, gnats and mosquitoes, dilute at the rate of 60 milliliters per 3.8L of water. REMOVE ANIMALS BEFORE SPRAYING. Apply as a fog or fine mist (at approximately 60 milliliters per 28 cu. meters of space), directing the spray toward the ceiling and upper corners, and not directly towards the animals or birds. For best results, close doors and windows before spraying, and keep them closed for 10 to 15 minutes. Applicator should vacate the treated area ventilate it prior to returning. Milking equipment should be covered before spraying and washed thoroughly before use. Repeat application as necessary.

STORAGE: Storage in a cool, dry place. Keep container closed.

DISPOSAL: Triple or pressure rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instruction for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Disposal of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Product Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use.

Medical 1-800-498-5743
? 1-800-621-8829

Manufactured By:
Chem-Tech, Ltd.
1006 North Highway 5
Pleasantville, IA 50225
United States

Prozap® is a Registered Trademark of Neogen Corporation.

Marketing Statements :

<Company Logo>
ChemTech, Ltd.*
***ChemTech, Ltd is a**
Wholly Owned
Subsidiary of Neogen
Corporation.

<Company Logo>
Neogen Corporation
Neogen Corporation

Code : (N.I.R.L.) NR.NA..

Item No.: 249279515-CAN
Net Contents: 32 ounce
UPC: 0 45446 02550 7
Format: UPC-A
Front Label: 9515LABF
Back Label: 9515LABB

Item No.: 249279515-CAN
Net Contents: 6 x 32 ounce
UPC: 000 45446 02555 2
Format: Interleaved 2-Of-5
Label: Case Label